

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# Covert Cockroach Gel Bait

**ACTIVE CONSTITUENT: 21.5 g/kg IMIDACLOPRID**

|       |    |             |
|-------|----|-------------|
| GROUP | 4A | INSECTICIDE |
|-------|----|-------------|


*For control of cockroaches in the situations specified in the Directions For Use.*

The syringe must not be sold separately.  
Do not destroy box while product still remains

Contents: 15g 30g 60g 120g (4 x 30g)

APVMA Approval No. 68912/59367

PCT INTERNATIONAL PTY LTD  
(PCT Holdings Pty Ltd ABN 11 099 023 962)  
1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>  
CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL  
HOURS) FREECALL 1800 630 877



Company: PCT Holdings Pty Ltd  
Product: **Covert Cockroach Gel Bait**  
Label Part: Outer box  
Text above this line is not part of the label

Page 2 of 4

Date: 23 Dec 2013

## DIRECTIONS FOR USE

Restraints

DO NOT place in areas that are routinely washed.

| SITUATION                                                                                                                                                                                                   | PEST                                                                        | RATE                                                                                                                                                                                                                                                    | CRITICAL COMMENTS                                                                                                                                                                                                                                                                                                                                                                                     |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Barracks, farm buildings, food processing establishments, factories, homes, hospitals, meatworks (non-product areas), offices, restaurants, schools, shops, store-rooms, transport vehicles and facilities. | <u>Small cockroaches</u> eg German cockroach ( <i>Blattella germanica</i> ) | <u>Light infestations</u><br>1 placement per m <sup>2</sup> .<br><br><u>Heavy infestations</u><br>2 placements per m <sup>2</sup> .<br><br>1 placement = a 5 mm ball (about the size of a small pea or approximately 0.1 g) or a line about 2.5 cm long | Apply gel in out of the way locations where cockroaches occur e.g., under fridges, near water heaters, behind microwave ovens, garbage bins etc. Re-assessment of treated areas one month after initial baiting is recommended to evaluate if additional baiting is required. Bait placements should be replenished when consumed or at least every three months as long as pest populations persist. |
|                                                                                                                                                                                                             | <u>Large cockroaches</u> eg American cockroach ( <i>Periplaneta</i> spp.)   | <u>Light infestations</u><br>2 placement per m <sup>2</sup> .<br><br><u>Heavy infestations</u><br>3 placements per m <sup>2</sup> .                                                                                                                     | <i>Periplaneta</i> spp. require more bait because of the quantity which can be consumed by a single cockroach. Reassess baiting effectiveness as above.                                                                                                                                                                                                                                               |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

Do not allow to contact food, livestock feed or their water. Do not place gel on surfaces where food is prepared or on Utensils that are used for food preparation, on food packing materials or on surfaces where food packing materials are stored.

## GENERAL INSTRUCTIONS

Apply using a suitable applicator for gel baits. Gel should be applied out of reach of children, pets and other animals. Do not place gel on surfaces which are regularly washed.

## **INSECTICIDE RESISTANCE WARNING**

For insecticide resistance management COVERT 

|              |           |                    |
|--------------|-----------|--------------------|
| <b>GROUP</b> | <b>4A</b> | <b>INSECTICIDE</b> |
|--------------|-----------|--------------------|

 COCKROACH GEL BAIT is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to COVERT COCKROACH GEL BAIT and other group 3A insecticide may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if COVERT COCKROACH GEL BAIT or other group 4A insecticides are used repeatedly. The effectiveness of COVERT COCKROACH GEL BAIT on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, PCT Holdings Pty Ltd accepts no liability for any losses that may result from the failure of COVERT COCKROACH GEL BAIT to control resistant insects. COVERT COCKROACH GEL BAIT may be subject to specific resistance management strategies. For further information, contact your local supplier, PCT Holdings Pty Ltd representatives or local agricultural department agronomist.

## **PROTECTION OF WILDLIFE FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with gel or used syringes.

## **STORAGE, SPILLAGE AND DISPOSAL**

**Keep out of reach of children.** Do not store in direct sunlight. Store in the closed, original container in a cool dry place. Completely empty syringe in normal use. Dispose of empty syringe and carton by wrapping in paper, placing in plastic bag and putting in the garbage.

## **SAFETY DIRECTIONS**

Will irritate the eyes. Avoid contact with eyes. Wash hands after use.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## **MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet (MSDS). A MSDS for COVERT COCKROACH GEL BAIT is available from PCT Holdings Pty Ltd on request.

## **WARRANTY**

PCT Holdings Pty Ltd makes no warranty expressed or implied, concerning the use of this product other than that indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# Covert Cockroach Gel Bait

**ACTIVE CONSTITUENT: 21.5 g/kg IMIDACLOPRID**

|       |           |             |
|-------|-----------|-------------|
| GROUP | <b>4A</b> | INSECTICIDE |
|-------|-----------|-------------|

*For control of cockroaches in the situations specified in the Directions For Use.*

BEFORE USING THIS PRODUCT READ ALL DIRECTIONS FOR USE ON OUTER  
DO NOT DESTROY OUTER PACK WHILE PRODUCT STILL REMAINS  
PACK. RETURN PLASTIC SYRINGE TO OUTER PACK AFTER USE.  
SALE OF THIS PLASTIC SYRINGE EXCEPT IN UNOPENED OUTER PACK IS ILLEGAL

## STORAGE, SPILLAGE AND DISPOSAL

**Keep out of reach of** children. Do not store in direct sunlight. Store in the closed, original container in a cool dry place. Completely empty syringe in normal use. Dispose of empty syringe and carton by wrapping in paper, placing in plastic bag and putting in the garbage.

## SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS). A MSDS for COVERT COCKROACH GEL BAIT is available from PCT Holdings Pty Ltd on request.

Net Contents: 15g 30g 60g

APVMA Approval No. 68912/59367

PCT INTERNATIONAL PTY LTD (PCT Holdings Pty Ltd ABN 11 099 023 962)  
1/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pct.au.com>  
CUSTOMER SERVICE FREECALL 1800 630 877  
EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

**Batch No.:**

**DOM:**